



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 213/00		A2	(11) International Publication Number: WO 99/15503
			(43) International Publication Date: 1 April 1999 (01.04.99)
(21) International Application Number: PCT/US98/19788 (22) International Filing Date: 22 September 1998 (22.09.98) (30) Priority Data: 60/060,680 25 September 1997 (25.09.97) US 9806419.9 25 March 1998 (25.03.98) GB (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DAVIES, Ian, W. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). GERENA, Linda [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). JOURNET, Michel [FR/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). LARSEN, Robert, D. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PYE, Philip, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ROSSEN, Kai [DE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: PROCESS FOR MAKING DIARYL PYRIDINES USEFUL AS COX-2 INHIBITORS			
<div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>			
(57) Abstract The invention encompasses a process for making compounds of Formula (I) useful in the treatment of cyclooxygenase-2 mediated diseases.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTIONPROCESS FOR MAKING DIARYL PYRIDINES USEFUL AS COX-2
INHIBITORS5 BACKGROUND OF THE INVENTION

This invention concerns a process for making certain anti-inflammatory compounds. In particular, the application concerns a process for making compounds of formula I as disclosed hereinunder, which compounds are potent cyclooxygenase-2 inhibitors.


10 Non-steroidal, antiinflammatory drugs exert most of their antiinflammatory, analgesic and antipyretic activity and inhibit hormone-induced uterine contractions and certain types of cancer growth through inhibition of prostaglandin G/H synthase, also known as cyclooxygenase. Up until recently, only one form of cyclooxygenase
15 had been characterized, this corresponding to cyclooxygenase-1 or the constitutive enzyme, as originally identified in bovine seminal vesicles. Recently the gene for a second inducible form of cyclooxygenase (cyclooxygenase-2) has been cloned, sequenced and characterized from chicken, murine and human sources. This enzyme is distinct from the
20 cyclooxygenase-1 which has now also been cloned, sequenced and characterized from sheep, murine and human sources. The second form of cyclooxygenase, cyclooxygenase-2, is rapidly and readily inducible by a number of agents including mitogens, endotoxin, hormones, cytokines and growth factors. As prostaglandins have both
25 physiological and pathological roles, we have concluded that the constitutive enzyme, cyclooxygenase-1, is responsible, in large part, for endogenous basal release of prostaglandins and hence is important in their physiological functions such as the maintenance of gastrointestinal integrity and renal blood flow. In contrast, we have
30 concluded that the inducible form, cyclooxygenase-2, is mainly responsible for the pathological effects of prostaglandins where rapid induction of the enzyme would occur in response to such agents as inflammatory agents, hormones, growth factors, and cytokines. Thus, a selective inhibitor of cyclooxygenase-2 will have similar
35 antiinflammatory, antipyretic and analgesic properties to a conventional

non-steroidal antiinflammatory drug, and in addition would inhibit hormone-induced uterine contractions and have potential anti-cancer effects, but will have a diminished ability to induce some of the mechanism-based side effects. In particular, such a compound should
5 have a reduced potential for gastrointestinal toxicity, a reduced potential for renal side effects, a reduced effect on bleeding times and possibly a lessened ability to induce asthma attacks in aspirin-sensitive asthmatic subjects.

WO 96/24585 published August 15, 1996 and WO 96/10012,
10 published April 4, 1996 disclose methods of making 2-aryl-3-aryl-pyridines. In the invention as disclosed hereinunder, 2-aryl-3-aryl-pyridines are prepared in a simple to conduct, 2 step conversion of a Weinreb amide to the penultimate ketosulfone from readily available starting materials. It is, therefore, surprisingly convenient and more
15 efficient than the previously described procedure, in which the 2-aryl-3-aryl pyridine was constructed by serial stepwise addition of the aryl groups to the central pyridine ring. Moreover, the process of the instant invention is also surprisingly superior in that expensive palladium reagents are not required nor is the cumbersome protection/de-
20 protection sequence of the prior art process.


SUMMARY OF THE INVENTION

The invention encompasses a process for making cyclooxygenase-2 intermediates such as the compound of structural
25 formula 5.



5

Compound 5 is useful in making potent cyclooxygenase-2 inhibitors of structural formula I, which are useful in the treatment of inflammation and other cyclooxygenase-2 mediated diseases



5 wherein:

R¹ is selected from the group consisting of

- (a) CH₃,
- (b) NH₂,
- (c) NHC(O)CF₃,
- 10 (d) NHCH₃;

Ar is a mono-, di-, or trisubstituted phenyl or pyridinyl (or the N-oxide thereof), wherein the substituents are chosen from the group consisting of

- (a) hydrogen,
- 15 (b) halo,
- (c) C₁₋₄alkoxy,
- (d) C₁₋₄alkylthio,
- (e) CN,
- (f) C₁₋₄alkyl,
- 20 (g) C₁₋₄fluoroalkyl, and


R² is chosen from the group consisting of

- (a) F, Cl, Br, I
- (b) CN,
- (c) azide.

DETAILED DESCRIPTION OF THE INVENTION

In a first aspect, the invention encompasses a process for making a compound of Formula 5, which is a COX-2 intermediate,


5



5

the process comprising:


10 reacting a compound of formula 13



13

wherein X is a halogen belonging to the group consisting of chlorine,
15 bromine and fluorine,


with a compound of formula 9



wherein Ar is a mono-, di-, or trisubstituted phenyl or pyridinyl (or the
 5 N-oxide thereof), wherein the substituents are chosen from the group
 consisting of

- (a) hydrogen,
- (b) halo,
- (c) C₁₋₄alkoxy,
- 10 (d) C₁₋₄alkylthio,
- (e) CN,
- (f) C₁₋₄alkyl,
- (g) C₁₋₄fluoroalkyl,

15 to yield a compound of formula 15




15

wherein Ar is described above,

20

and oxidizing the compound of formula 15 using an oxidizing agent, and
 optionally a catalyst and an acid to yield a compound of formula 5.

In a second aspect, the invention encompasses a process for
 making a compound of formula 13




13

wherein X is a halogen belonging to the group consisting of chlorine, bromine and fluorine,

5

comprising reacting a compound of formula 12




12

with magnesium in the presence of a solvent/co-solvent mixture to yield a compound of formula 13.

10

In a third aspect, the invention encompasses a process for making a compound of formula I



I

wherein:

R¹ is selected from the group consisting of

- (a) CH₃,
- (b) NH₂,
- (c) NHC(O)CF₃,
- (d) NHCH₃;

5

Ar is a mono-, di-, or trisubstituted phenyl or pyridinyl (or the N-oxide thereof), wherein the substituents are chosen from the group consisting of

- (a) hydrogen,
- (b) halo,
- (c) C₁₋₄alkoxy,
- (d) C₁₋₄alkylthio,
- (e) CN,
- (f) C₁₋₄alkyl,
- (g) C₁₋₄fluoroalkyl, and

10


15

R² is chosen from the group consisting of

- (a) F, Cl, Br, I
- (b) CN,
- (c) azide;

20

comprising reacting a compound of formula 12




12

wherein X is a halogen belonging to the group consisting of iodine, chlorine, bromine and fluorine,


25

with magnesium in the presence of a solvent/co-solvent mixture to yield a compound of formula 13

8




reacting the compound of formula 13 with a compound of formula 9



5

wherein Ar is described above,


10 to yield a compound of formula 15



wherein Ar is described above,

15


and oxidizing the compound of formula 15 using an oxidizing agent and optionally a catalyst under acid conditions to yield a compound of formula




5

5 wherein R1 is described above,

condensing a compound of formula A1 or A2



A1




A2

10


under acidic conditions, optionally in the presence of a non-reactive solvent and in the presence of an ammonium reagent, with compound 5 to yield a compound of Formula I.

15 In a fourth aspect of the invention, the process encompasses a method for making a compound of formula II:



II


comprising reacting a compound of formula 12



12

wherein X is a halogen belonging to the group consisting of iodide,
5 chloride, bromide and fluoride;


with magnesium in the presence of a solvent/co-solvent mixture to yield
a compound of formula 13



13

10


reacting the compound of formula 13 with a compound of formula 9a




9a

15

to yield a compound of formula 15a



**15a**

- 5 and oxidizing the compound of formula **15a** using an oxidizing agent and optionally a catalyst under acid conditions to yield a compound of formula

**5a**

10

condensing a compound of formula **A1** or **A2**

**A1****A2**

- 15 wherein R^2 is F, Cl, Br, I, CN, or azide;

under acidic conditions, and optionally in the presence of a non-reactive solvent and in the presence of an ammonium reagent, with compound 5a

5 to yield a compound of Formula II.

A further aspect of this invention is realized when A1 is employed wherein the acidic condition consists of the addition of acetic or propionic acid, the non-reactive solvent is tetrahydrofuran, dioxane, C1-6alkanol, chlorobenzene, dichlorobenze, xylene or toluene and the
10 ammonium reagent is ammonia, ammonium acetate or ammonium propionate.

A further aspect of this invention is realized when A2 is employed wherein the acidic condition consists of the addition of acetic acid, methanesulfonic acid or propionic acid or a mixture thereof,
15 preferably a mixture of propionic acid and methanesulfonic acid, the non-reactive solvent is tetrahydrofuran, dioxane, C1-6alkanol, chlorobenzene, dichlorobenze, xylene or toluene and the ammonium reagent is ammonia, ammonium acetate, ammonium hydroxide and ammonium propionate, preferably ammonium hydroxide.

20 For purposes of this specification, the reactions, unless otherwise specified, are generally carried out in a solvent such as benzene, chlorobenzene, dichlorobenze, toluene and xylene; ethereal solvents such as diethyl ether, di-n-butyl and diisopentyl ethers, anisole, cyclic ethers such as tetrahydropyran, 4-methyl-1,3-dioxane,
25 dihydropyran, tetrahydrofurfuryl, methyl ether, ethyl ether, 2-ethoxytetrahydrofuran and tetrahydrofuran (THF); ester solvents including ethyl and isopropyl acetate; halo carbon solvents including mono or dihalo C1-4 alkyl such as dichloromethane; C6-10 linear, branched or cyclic hydrocarbon solvents including hexane; and nitrogen
30 containing solvents including N,N-dimethylacetamide, N,N-dimethylformamide (DMF), N-ethylpyrrolidinone, N-methylpyrrolidinone, and acetonitrile. Preferable solvents are alcohol, dichloromethane, THF and DMF.

For purposes of this specification X is a halogen belonging to the group consisting of iodide, chloride, bromide or fluoride, preferably chloride and compound 12 is commercially available.

Regarding the first aspect of the invention, the oxidizing agent belongs to the group consisting of hydrogen peroxide, oxone, hydrogen peroxide/acetic acid and the like, preferably oxone or hydrogen peroxide and the catalyst is Na_2WO_4 . The acid shall include acetic, propionic or another carboxylic acid, hydrochloric acid or sulfuric acid and the like. The pH is maintained at about 2 to about 5, preferably about 2-4. The reaction is preferably conducted employing an acid such as sulfuric acid.

The molar ratio of 13 to 9 can typically be varied from about 1:1 to about 2:1; preferably about 1:5 to about 1. Excess compound 13 relative to compound 9 is typically used. The molar ratio of compound 15 to oxidizing agent can typically be varied from about 1:1 to about 1:10. The molar ratio of the catalyst to compound 15 can typically be varied from about 1:1 to about 1:1000, preferably about 1:100. The reaction may conveniently be conducted at a temperature range of about 0 to about 100°C; preferably about 50 to about 60°C and is allowed to proceed until substantially complete in from 1 to 24 hours.

For purposes of this specification, in the second aspect of the invention the solvent/co-solvent mixture shall include solvent mixtures such as toluene/tetrahydrofuran, tetrahydrofuran/diethylether, toluene/diethylether, tetrahydrofuran/methyl-t-butylether, toluene/methyl-t-butylether, toluene/dioxane, tetrahydrofuran/dioxane and the like, preferably toluene/tetrahydrofuran.

The molar ratio of compound 12 to solvent/co-solvent mixture can typically be varied from about 1:20 to about 1:1, preferably about 1:3 to about 1:1. The molar ratio of solvent to co-solvent can typically be varied from about 0.5:4 to about 1:1, preferably about 1:2 to about 1:1. The molar ratio of compound 12 to magnesium can typically be varied from about 1:2 to about 1:1. Generally, compound 12 is mixed with the required amount of co-solvent and added to the solvent containing magnesium and the reaction is conveniently conducted at a temperature range of about 0 to about 40°C; preferably about 10 to about

35°C. The reaction is allowed to proceed until substantially complete in from 1 to 5 hours; typically 1 to 2 hours.

Regarding the third aspect of the invention, as will be appreciated by those of skill in the art, in the general case the reagents themselves provide the acidic condition. Therefore, the use of a non-reagent acid is not necessary. However, the addition of an acid, such as acetic or propionic or another carboxylic acid or a mixture of acids such as propionic acid and methanesulfonic acid are within the scope of the invention.

For purposes of this specification non-reactive solvent includes tetrahydrofuran, dioxane, C1-6alkanol, chlorobenzene, dichlorobenze, xylene and toluene.

For purposes of this specification, the ammonium reagent is intended to include ammonia, ammonium salts such as ammonium acetate and ammonium propionate and aqueous ammonia such as ammonium hydroxide. Moreover a mixture ammonia reagent species is included in the term ammonia reagent.

The molar ratio of compound A1 or A2 to 5 can typically be varied from about 3:1 to about 1:2; preferably about 1:1 to about 1.5. Excess compound A1 is typically used. The molar ratio of compound A1 or A2 to ammonium reagent can typically be varied from about 1:1 to about 1:10. The reaction step may conveniently be conducted at a temperature range of about 40 to about 180°C; preferably about 80 to about 140°C and is allowed to proceed until substantially complete in from about 2 to about 18 hours; typically about 6 to about 12 hours.


With regard to the third aspect of the invention, R² is preferably halogen, most preferably F, Br, or Cl, most preferably Cl.

With regard to all aspects of the invention a preferred sub-genus of formula I is that wherein Ar is a mono-, or disubstituted pyridinyl. Within this sub-genus, the 3-pyridinyl isomers are particularly preferred.

Again with regard to all aspects of the invention another preferred sub-genus of formula I is that wherein R¹ is CH₃ or NH₂. Generally, CH₃ is preferred for COX-2 specificity and NH₂ is preferred for potency.

Again with regard to all aspects of the invention another preferred sub-genus of formula I is that wherein the Ar is unsubstituted or substituted with CH₃.

In a fifth aspect of the invention the compounds of formula
5 A2 are described:



A2

wherein R² is:

- (a) halogen
- 10 (b) CN,
- (c) azide
- (d) C₂₋₆ alkyl optionally substituted with 1 to 3 groups of C₁₋₆ alkyl, hydroxy, halogen, carbonyl, CO₂, NO₂, OC₁₋₆ alkyl; SC₁₋₆ alkyl, N(C₁₋₆ alkyl)₂
- 15 (e) C₆₋₁₀ aryl or heteroaryl optionally substituted with 1 to 3 groups of C₁₋₆ alkyl, hydroxy, halogen, carbonyl, CO₂, NO₂, OC₁₋₆ alkyl; SC₁₋₆ alkyl, N(C₁₋₆ alkyl)₂.

A preferred compound is realized when R² is halogen, alkyl, phenyl or substituted phenyl. Most preferably compounds are
20 realized when R² is fluorine, bromine, iodine, chlorine, ethyl, isopropyl, phenyl, trifluorophenyl

As used herein, "alkyl" is intended to include branched, cyclic and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Examples of cycloalkyl
25 groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

As used herein, "aryl" is intended to include aryls and heteroaryls, both substituted and unsubstituted, which are defined as carbazolyl, furyl, thienyl, pyrrolyl, isothiazolyl, imidazolyl, isoxazolyl, thiazolyl, oxazolyl, pyrazolyl, pyrazinyl, pyridyl, pyrimidyl, purinyl or
30 quinolinyl as well as aromatic rings e.g., phenyl, substituted phenyl and like groups as well as rings which are fused, e.g., naphthyl and the like.

Substitution can be 1 to 3 groups of C₁₋₆ alkyl, hydroxy, halogen, carbonyl, CO₂, NO₂, OC₁₋₆ alkyl; SC₁₋₆ alkyl; N(C₁₋₆ alkyl)₂ and the like.

As used herein, "halogen" is intended to include chlorine, fluorine, bromine and iodine.

- 5 The Compound of Formula I is useful for the relief of pain, fever and inflammation of a variety of conditions including rheumatic fever, symptoms associated with influenza or other viral infections, common cold, low back and neck pain, dysmenorrhea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, arthritis,
10 including rheumatoid arthritis degenerative joint diseases (osteoarthritis), gout and ankylosing spondylitis, bursitis, burns, injuries, following surgical and dental procedures. In addition, such a compound may inhibit cellular neoplastic transformations and metastatic tumor growth and hence can be used in the treatment of cancer.
- 15 Compounds of formula I may also be useful for the treatment of dementia including pre-senile and senile dementia, and in particular, dementia associated with Alzheimer Disease (ie Alzheimer's dementia).

- By virtue of its high cyclooxygenase-2 (COX-2) activity and/or its selectivity for cyclooxygenase-2 over cyclooxygenase-1 (COX-1)
20 as defined above, compounds of formula I will prove useful as an alternative to conventional non-steroidal antiinflammatory drugs (NSAID'S) particularly where such non-steroidal antiinflammatory drugs may be contra-indicated such as in patients with peptic ulcers, gastritis, regional enteritis, ulcerative colitis, diverticulitis or with a
25 recurrent history of gastrointestinal lesions; GI bleeding, coagulation disorders including anemia such as hypoprothrombinemia, haemophilia or other bleeding problems (including those relating to reduced or impaired platelet function); kidney disease (eg impaired renal function); those prior to surgery or taking anticoagulants; and
30 those susceptible to NSAID induced asthma.

Compounds of the present invention are inhibitors of cyclooxygenase-2 and are thereby useful in the treatment of cyclooxygenase-2 mediated diseases as enumerated above. This activity is illustrated by their ability to selectively inhibit cyclooxygenase-2 over

cyclooxygenase-1. Accordingly, in one assay, the ability of the compounds of this invention to treat cyclooxygenase mediated diseases can be demonstrated by measuring the amount of prostaglandin E₂ (PGE₂) synthesized in the presence of arachidonic acid, cyclooxygenase-1 or cyclooxygenase-2 and a compound of formula I. The IC₅₀ values represent the concentration of inhibitor required to return PGE₂ synthesis to 50 % of that obtained as compared to the uninhibited control. Illustrating this aspect, we have found that the Compounds of the Examples are more than 100 times more effective in inhibiting COX-2 than they are at inhibiting COX-1. In addition they all have a COX-2 IC₅₀ of 1 nM to 1 mM. By way of comparison, Ibuprofen has an IC₅₀ for COX-2 of 1 mM, and Indomethacin has an IC₅₀ for COX-2 of approximately 100 nM.

For the treatment of any of these cyclooxygenase mediated diseases, compounds of formula I may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle sheep, dogs, cats, etc., the compound of the invention is effective in the treatment of humans.

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) all operations were carried out at room or ambient temperature, that is, at a temperature in the range 18-25°C; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 pascals: 4.5-30 mm. Hg) with a bath temperature of up to 60°C; the course of reactions was followed by thin layer chromatography (TLC) or High Pressure Liquid Chromatography (HPLC) and reaction times are given for illustration only; melting points are uncorrected and 'd' indicates decomposition; the melting points given are those obtained for the materials prepared as described; polymorphism may result in isolation of materials with different

- melting points in some preparations; the structure and purity of all final products were assured by at least one of the following techniques: TLC, mass spectrometry, nuclear magnetic resonance (NMR) spectrometry or microanalytical data; yields are given for illustration only; when given,
- 5 NMR data is in the form of delta (δ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, determined at 300 MHz or 400 MHz using the indicated solvent; conventional abbreviations used for signal shape are: s. singlet; d. doublet; t. triplet; m. multiplet; br. broad; etc.: in addition
- 10 "Ar" signifies an aromatic signal; chemical symbols have their usual meanings; the following abbreviations have also been used v (volume), w (weight), b.p. (boiling point), m.p. (melting point), L (liter(s)), mL (milliliters), g (gram(s)), mg (milligrams(s)), mol (moles), mmol (millimoles), eq (equivalent(s)).

15


The following abbreviations have the indicated meanings:

Alkyl Group Abbreviations

Me	=	methyl
Et	=	ethyl
n-Pr	=	normal propyl
i-Pr	=	isopropyl
n-Bu	=	normal butyl
i-Bu	=	isobutyl
s-Bu	=	secondary butyl
t-Bu	=	tertiary butyl
c-Pr	=	cyclopropyl
c-Bu	=	cyclobutyl
c-Pen	=	cyclopentyl
c-Hex	=	cyclohexyl

EXAMPLE 1

- 20 5-Chloro-3(methylsulfonyl)phenyl-2-(3-pyridyl)-pyridine; Compound 1



3-Amino-2-chloroacrolein	140g (1.31 mol)
Ketosulfone	136g (4.69 mol)
Methanesulfonic acid	99.2g (10.3 mol)
<i>n</i> -Propionic acid	690 mL (92.5 mol)
Ammonium hydroxide (14.8M)	600 mL (88.8 mol)
Toluene	1.35 L


10

A mixture of *n*-propionic acid (400 mL), 3-amino-2-chloroacrolein (140 g, 1.31 mol), ketosulfone (136 g, 0.469 mol), toluene (1.35 L), propionic acid (690 mL, 92.5 mol), methanesulfonic acid (67mL, 10.3 mol) was heated to reflux (114°C) for 12 hours with the azeotropic removal of water. The reaction solution was cooled to ambient temperature and diluted with isopropyl acetate (1 L). Water (1 L) was added and the aqueous phase was neutralized with concentrated ammonium hydroxide solution (600 mL). The organic layer was washed with a 1:1 mixture of brine/water (2 X 1 L) and water (1 L). The combined aqueous layers were extracted with isopropyl acetate (900 mL). The combined organic layers were treated with Darco G-60 then concentrated. Recrystallization from isopropylacetate/ hexanes gave a colorless solid (123.1g, 65%) mp 135 °C (DSC).

25


EXAMPLE 1A

5-Chloro-3(methylsulfonyl)phenyl-2-(3-pyridyl)-pyridine; Compound 1




5	2-Chloromalondialdehyde	4.8 g (0.045 mol)
	Ketone B	5.0 g (0.018 mol)
	Propionic acid	30 mL
	Ammonium Acetate	8.4 g (0.11 mol)

A mixture of ketone B (5.0 g), 2-chloromalondialdehyde (4.8 g) and ammonium acetate were heated to 130°C. The acetic acid produced was removed by distillation and heating continued at 136°C for 15 hours. The reaction mixture was basified with sodium carbonate, water was added and the product was extracted into dichloromethane (2 x 150 mL). The organic layers were carbon treated (Dowex), dried (MgSO₄) and the solvent removed to afford 1 as an off white solid (3.4 g, 55% yield).



20	2-Chloromalondialdehyde	220 mg (2.1 mmol)
	Oxalyl Chloride	180mL (2.1 mmol)
	Toluene	3 mL
	N,N-Dimethylformamide	20mL


N,N-dimethyl formamide was added to a slurry of 2-chloromalondialdehyde (220 mg) in toluene. Oxalyl chloride was added and the reaction mixture was stirred until complete dissolution occurred.



	Ketone B	500 mg (1.8 mmol)
	Lithium bis(trimethylsilyl)amide (1 M in THF)	1.8 mL (1.8 mmol)
5	Tetrahydrofuran	15 mL
	2,3-Dichloroacrolein in toluene	2.1 mmol in 3 mL toluene
	Ammonium acetate	1.0 g

Lithium bis(trimethylsilyl)amide (1.8 mL; 1 M in THF) was added dropwise to ketone B (500 mg) in THF (15 mL) at -78°C. The reaction mixture was warmed to ambient temperature for 1 hour to form the lithium enolate of B (see the generic formula B1) before recooling to -78°C. A solution of 2,3-dichloroacrolein was added and the temperature allowed to warm to room temperature. After 1 hour ammonia gas was passed through the solution and after 30 minutes ammonium acetate (1 g) was added. The reaction mixture was warmed to 60°C for 1 hour and poured into aqueous sodium hydroxide (2 M; 100 mL). The product was extracted with dichloromethane (2 x 150 mL), dried (MgSO₄) and the solvent removed to afford 1 (500 mg; 80%).

20

EXAMPLE 2**Weinreb**

25

Methyl 6-methylnicotinate
 N,O-Dimethylhydroxylamine


21.56 g (0.143 mol)
 13.9 g (0.229 mol)

Tetrahydrofuran 150 mL
 Isopropylmagnesium chloride 110 mL (0.220 mol)
 (2.0M in THF)
 Toluene 180 ml

5

A solution of Methyl 6-methylnicotinate (21.56 g), and *N,O*-dimethylhydroxylamine (13.9 g) in THF (150 mL) was cooled to -10°C. Isopropylmagnesium chloride (110 mL) was added over 2.5 h. The reaction mixture was poured into aqueous acetic acid (10 vol%, 126 mL) at 5 °C. Toluene (60 mL) was added to the mixture, then the layers were separated. The aqueous layer was extracted with toluene (2 x 60 mL) and the solvent removed. Solid impurities were removed by filtration and the filtrate was concentrated to afford the Weinreb amide as a light orange oil (24.2 g, 94.3%).

15

EXAMPLE 3**Grignard**

20

4-thiomethylbenzyl chloride 566g (3.28 mol)
 Magnesium 191g (7.86 mol)
 Toluene 9L
 Tetrahydrofuran 0.545L
 Weinreb amide 2 300g (1.66 mol)


25

A mixture of magnesium (191g, 7.86 mol) toluene (4L), 4-thiomethylbenzyl chloride (566g, 3.28 mol) and tetrahydrofuran (0.545L, 6.73 mol) were charged over 3 - 4 hours. An additional flask was charged with Weinreb amide (300g, 1.66 mol) and toluene (1.7L) and cooled to -20 °C. The Grignard solution prepared above was added over 30 minutes and the mixture was aged for 1 hour. The reaction mixture

30

was quenched by the addition of 50% aqueous acetic acid (0.5L). Toluene (1L) and water (1L) were added and the layers were separated. The aqueous layer was extracted with toluene (2 x 2L). The combined organic extracts were extracted with dilute hydrochloric acid (1 x 2L). Ethyl acetate was added to the aqueous layer and the pH was adjusted with ammonia (0.6L). The phases were separated and the aqueous layer was extracted with ethyl acetate (2x 1.25L). The combined extracts were concentrated on a rotary evaporator to give a light yellow solid (326.5g, 76%)

10


EXAMPLE 4Oxidation

15

Ketosulfide	270g (1.05 mol)
Methanol	2.70L
20 Sodium tungstate	6.0g (0.02 mol)
Water	5.20L
Sulfuric Acid (2N)	0.02L
Hydrogen peroxide (30%)	380 mL (3.0 mol)

25 A mixture of ketosulfide (270g, 1.05 mol), sulfuric acid (2N) (20 mL), and methanol (2.70L) was heated at 55 °C. An aqueous solution of sodium tungstate (6.0g, 0.02 mol) was added then hydrogen peroxide (380 mL) was added over 1 hour. Water (3L) was added and the mixture was cooled to ambient temperature then filtered. The solids were washed with water (2L) and dried under vacuum with a stream of nitrogen to give ketosulfone 8 (250.2g, 82.5%) as a colorless solid.

30

EXAMPLE 5

5

Chloromalonaldol 400 g (3.76 mol)
 Aqueous ammonia (30%, 14.8 N) 370 mL (5.48 mol)
 Isopropyl alcohol 6.4 L

10


To a flask was charged with chloromalonaldol (400 g, 3.76 mol), and isopropyl alcohol (400 mL). The solution was concentrated under reduced pressure with a continuous, slow feed of isopropyl alcohol (4.0 L total). The resulting dark brown liquid was diluted with isopropyl alcohol (400 mL). The mixture was added to a cooled (5 °C) solution of 30% aqueous ammonia (370 mL) in isopropyl alcohol (2 L). The mixture was aged for 3 hours and the product was collected by filtration (373 g, 93%)

15

EXAMPLE 5A

20

A number of routes are available for the preparation of chloromalonaldialdehyde.


Preparation from 1,1,2,3,3-Pentachloropropane

25

A detailed experimental is published in Houben-Weyl-Muller: Methoden der Organischen Chemie, 4th Edit., Vol 7/1, Thieme Verlag, Stuttgart, 1954, page 119. The starting material 1,1,2,3,3-pentachloropropane is commercially available from Pfaltz and Bauer.

5

Preparation from Mucochloric Acid



- 10 The following is a slight variation of the original procedure of Dieckmann (*Ber. Deut. Chem. Ges.* 1904, 37, 4638).

15	Mucochloric acid Aniline Water	50.0 g (0.30 mol) 54 mL (0.60 mol) 1000 mL
----	--------------------------------------	--

- 20 To a solution of aniline in water at 85°C in a vigorously stirred 2 L flask was added mucochloric acid in small portions over 30 min. On addition of the mucochloric acid, a yellow color develops, which quickly dissipated. The reaction mixture stayed heterogeneous and filtration of an aliquot after 30 min heating indicated completion of the reaction.

- 25 The reaction mixture was heated at 90°C for 60 min., cooled to 50°C and filtered. The filtercake was washed with 50 mL of 2N HCl and 100 mL of H_2O . The product was dried in a N_2 stream to give 57 g

(100% yield) of 3-anilido-2-chloro-acrolein as a gray solid. ^{13}C NMR (D_6 -DMSO in ppm): 108, 117, 124, 129, 140, 147, 182.

3-Anilido-2-chloro-acrolein 57 g (0.30 mol)
5 5N NaOH solution 120 mL (0.6 mol)


A solution of 3-anilido-2-chloro-acrolein in 120 mL of 5N NaOH was heated to 100°C for 90 min. The dark black solution was extracted twice with 50 mL each of MTBE.

10 The first organic wash removed most of the dark color from the solution, and the second organic wash was only lightly colored.

On cooling the aqueous phase, a crystalline precipitate formed. This product was the 3-chloromalondialdehyde Na salt.

15 The aqueous phase was acidified by the addition of 60 mL of 37% HCl solution. The aqueous phase was extracted (MTBE/THF 50/50, 400 mL total) and the combined organic phases were dried over MgSO_4 . After treatment with Darco G60 and filtration through a plug of SiO_2 , the solution was evaporated to give 19.6 g (62% overall yield) of chloromalondialdehyde as a dark solid. Recrystallization from ca. 10
20 mL of MTBE gave 11.13 g of pure chloromalondialdehyde as a tan solid.
 ^{13}C NMR (D_6 -DMSO in ppm): 113, 175 (broad).

Preparation from Chloroacetylchloride




Arnold (*Collect. Czech. Chem. Commun.* **1961**, 26, 3051) mentions the formation of 3-dimethylamino-2-chloro-acrolein by reaction of chloroacetic acid with the Vilsmeier reagent derived from POCl3 and DMF. A variation and extension of his procedure prepares chloromalonaldialdehyde as its Na salt.

Oxalylchloride (280 mL, 3.2 mol) was added at 10°C to 1000 mL of DMF. The reaction was highly exothermic and a heavy precipitate formed. After a 2 h age, chloroacetylchloride (110 mL, 1.4 mol) was added and the reaction mixture was warmed to 75°C for 3 hours. Analysis of an aliquot by ^1H NMR indicated complete consumption of the chloroacetylchloride and the reaction mixture was quenched by addition into 1 L of H2O. To the cooled solution was added 500 mL of a 50% NaOH solution. The reaction mixture is heated to reflux for 5 hours. On cooling a precipitate formed, which was filtered and washed with water. The tan solid was dried in a N2 stream to give 84 g of a tan solid (54% yield).

WHAT IS CLAIMED

1. A process for making compounds of Formula 5



R^1 is selected from the group consisting of

- (a) CH_3 ,
- (b) NH_2 ,
- (c) $NHC(O)CF_3$,
- (d) $NHCH_3$; and


Ar is a mono-, di-, or trisubstituted phenyl or pyridinyl (or the N-oxide thereof), wherein the substituents are chosen from the group consisting of

- (a) hydrogen,
- (b) halo,
- (c) C_{1-4} alkoxy,
- (d) C_{1-4} alkylthio,
- (e) CN,
- (f) C_{1-4} alkyl,
- (g) C_{1-4} fluoroalkyl,

the process comprising:

reacting a compound of formula 13


29

**13**

wherein X is a halogen belonging to the group consisting of iodine, chlorine, bromine and fluorine,


5

with a compound of formula 9

**9**

10

to yield a compound of formula 15

**15**

15 and oxidizing the compound of formula 15 using an oxidizing agent, and optionally a catalyst and an acid to yield a compound of formula 5.

2. A process according to Claim 1 wherein the oxidizing agent belongs to the group consisting of hydrogen peroxide, oxone, and hydrogen peroxide/acetic acid and the acid is acetic, propionic, methanesulfonic acid or sulfuric acid.

5

3. A process according to Claim 1 wherein the catalyst is Na_2WO_4 .

4. A process according to Claim 2 wherein the oxidizing agent is oxone or hydrogen peroxide and the acid is sulfuric acid.

10

5. A process according to Claim 1 wherein Ar is a mono- or di-trisubstituted 3-pyridinyl.

15

6. A process according to Claim 1 wherein R^1 is CH_3 or NH_2 .


7. A process according to Claim 1 wherein Ar is a mono- or di-substituted 3-pyridinyl and the substituents are selected from the group consisting of

20

- (a) hydrogen,
- (b) halo,
- (c) C_1 -3alkoxy,
- (d) C_1 -3alkylthio,
- (e) C_1 -3alkyl,
- (f) CF_3 , and
- (g) CN.

25


8. A process for making a compound of formula 13



13

wherein X is a halogen belonging to the group consisting of iodide,
5 chloride, bromide and fluoride;

comprising reacting a compound of formula 12



12

10

with magnesium in the presence of a solvent/co-solvent mixture to yield
a compound of formula 13.

15

9. A process according to Claim 8 wherein X is chloride.

10. A process according to Claim 8 wherein the
solvent/co-solvent mixture is selected from the group consisting of
toluene/tetrahydrofuran, tetrahydrofuran/diethylether,
20 toluene/diethylether, tetrahydrofuran/methyl-t-butylether,


toluene/methyl-t-butylether, toluene/dioxane and tetrahydrofuran/dioxane.

11. A process according to Claim 10 wherein the solvent/co-solvent mixture is toluene/tetrahydrofuran.

12. A process according to Claim 8 wherein the molar ratio of solvent to co-solvent is from about 0.5:4 to about 1:1.

13. A process according to Claim 12 wherein the molar ratio of solvent to co-solvent is from about 1:2 to about 1:1.

14. A process for making a compound of formula I



I

wherein:

R¹ is selected from the group consisting of

- (a) CH₃,
- (b) NH₂,
- (c) NHC(O)CF₃,
- (d) NHCH₃;

Ar is a mono-, di-, or trisubstituted phenyl or pyridinyl (or the N-oxide thereof), wherein the substituents are chosen from the group consisting of

- (a) hydrogen,
- (b) halo,
- (c) C₁₋₄alkoxy,
- (d) C₁₋₄alkylthio,

- (e) CN,
(f) C₁₋₄alkyl,
(g) C₁₋₄fluoroalkyl, and


R² is chosen from the group consisting of

5

- (a) F, Cl, Br, I
(b) CN,
(c) azide

comprising reacting a compound of formula 12

10




12

wherein X is a halogen belonging to the group consisting of iodide, chloride, bromide and fluoride;

15


with magnesium in the presence of a solvent/co-solvent mixture to yield a compound of formula 13




13

20


reacting the compound of formula 13 with a compound of formula 9




to yield a compound of formula 15




and oxidizing the compound of formula 15 using an oxidizing agent and optionally a catalyst under acid conditions to yield a compound of formula



condensing a compound of formula A1 or A2



A1



A2

- under acidic conditions, and optionally in the presence of a non-reactive solvent and in the presence of an ammonium reagent, with compound 5 to yield a compound of Formula I.

15. A process according to Claim 14 wherein the oxidizing agent belongs to the group consisting of hydrogen peroxide, oxone, and hydrogen peroxide/acetic acid and the acid is acetic, propionic, methanesulfonic acid or sulfuric acid.
16. A process according to Claim 14 wherein the catalyst is Na_2WO_4 .
17. A process according to Claim 15 wherein the oxidizing agent is oxone or hydrogen peroxide and the acid is sulfuric acid.
18. A process according to Claim 14 wherein Ar is a mono- or di-trisubstituted 3-pyridinyl.
19. A process according to Claim 1 wherein R^1 is CH_3 or NH_2 and R^2 is F, Br or Cl.
20. A process according to Claim 14 wherein Ar is a mono- or di-substituted 3-pyridinyl and the substituents are selected from the group consisting of
 - (a) hydrogen,
 - (b) halo,
 - (c) C_{1-3} alkoxy,

- (d) C₁-3alkylthio,
- (e) C₁-3alkyl,
- (f) CF₃, and
- (g) CN.

5

21. A process according to Claim 14 wherein X is chloride.

22. A process according to Claim 14 wherein the solvent/co-solvent mixture is selected from the group consisting of toluene/tetrahydrofuran, tetrahydrofuran/diethylether, toluene/diethylether, tetrahydrofuran/methyl-t-butylether, toluene/methyl-t-butylether, toluene/dioxane and tetrahydrofuran/dioxane.

15

23. A process according to Claim 22 wherein the solvent/co-solvent mixture is toluene/tetrahydrofuran.

24. A process according to Claim 14 wherein the molar ratio of solvent to co-solvent is from about 0.5:4 to about 1:1.

20

25. A process according to Claim 24 wherein the molar ratio of solvent to co-solvent is from about 1:2 to about 1:1.

25

26. A process according to Claim 14 employing A1 wherein the acidic condition consists of the addition of acetic or propionic acid, the non-reactive solvent is tetrahydrofuran, dioxane, C₁-6alkanol, or toluene and the ammonium reagent is ammonia, ammonium acetate and ammonium propionate.


30

27. A process according to 14 employing A2 wherein the acidic condition consists of the addition of acetic acid, methanesulfonic acid or propionic acid or a mixture thereof, the non-reactive solvent is tetrahydrofuran, dioxane, C₁-6alkanol, or toluene and the ammonium

reagent is ammonia, ammonium acetate, ammonium hydroxide and ammonium propionate.


28. A process according to 27 wherein the acidic condition consists of the addition of a mixture of propionic acid and methanesulfonic acid and the ammonium reagent is ammonium hydroxide.

29. A process for making a compound of formula II



II

comprising reacting a compound of formula 12




12


- wherein X is a halogen belonging to the group consisting of iodide, chloride, bromide and fluoride;

with magnesium in the presence of a solvent/co-solvent mixture to yield a compound of formula 13

38




reacting the compound of formula 13 with a compound of formula 9a




5

10 to yield a compound of formula 15a



15a


and oxidizing the compound of formula 15a using an oxidizing agent and optionally a catalyst under acid conditions to yield a compound of formula



5


5a

condensing a compound of formula A1 or A2



10

A1



A2

wherein R² is F, Cl, Br, I, CN, or azide;

under acidic conditions, and optionally in the presence of a non-reactive solvent and in the presence of an ammonium reagent, with compound 5a to yield a compound of Formula II.


30. A process according to Claim 29 wherein X is chloride and R² is Cl.

31. A process according to Claim 29 employing A1 wherein the acidic condition consists of the addition of acetic or

propionic acid, the non-reactive solvent is tetrahydrofuran, dioxane, C₁-6alkanol, or toluene and the ammonium reagent is ammonia, ammonium acetate and ammonium propionate.

- 5 32. A process according to 29 employing A2 wherein the acidic condition consists of the addition of acetic acid, methanesulfonic acid or propionic acid or a mixture thereof, the non-reactive solvent is tetrahydrofuran, dioxane, C₁-6alkanol, or toluene and the ammonium reagent is ammonia, ammonium acetate, ammonium hydroxide and
10 ammonium propionate.

33. A compound of structural formula A2



15

wherein R² is:

- (a) halogen
 (b) CN,
 (c) azide
 20 (d) C₂₋₆ alkyl optionally substituted with 1 to 3 groups of C₁-6 alkyl, hydroxy, halogen, carbonyl, CO₂, NO₂, OC₁-6 alkyl, SC₁-6 alkyl, N(C₁-6 alkyl)₂
 (e) C₆₋₁₀ aryl or heteroaryl optionally substituted with 1 to 3
 groups of C₁-6 alkyl, hydroxy, halogen, carbonyl, CO₂, NO₂, OC₁-6 alkyl;
 25 SC₁-6 alkyl, N(C₁-6 alkyl)₂.


34. A compound according to Claim 29 wherein R² is fluorine, bromine, iodine, chlorine, ethyl, isopropyl, phenyl, trifluorophenyl.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 213/61, C07C 223/02		A3	(11) International Publication Number: WO 99/15503
			(43) International Publication Date: 1 April 1999 (01.04.99)
(21) International Application Number: PCT/US98/19788 (22) International Filing Date: 22 September 1998 (22.09.98) (30) Priority Data: 60/060,680 25 September 1997 (25.09.97) US 9806419.9 25 March 1998 (25.03.98) GB (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DAVIES, Ian, W. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). GERENA, Linda [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). JOURNET, Michel [FR/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). LARSEN, Robert, D. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PYE, Philip, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ROSSEN, Kai [DE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 20 May 1999 (20.05.99)	

(54) Title: PROCESS FOR MAKING DIARYL PYRIDINES USEFUL AS COX-2 INHIBITORS



(57) Abstract

The invention encompasses a process for making compounds of Formula (I) useful in the treatment of cyclooxygenase-2 mediated diseases.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/19788

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D213/61 C07C223/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 24584 A (SEARLE & CO ;WEIER RICHARD M (US); LEE LEN F (US); PARTIS RICHARD) 15 August 1996 see page 50, scheme VII ----	14-18, 20-32
A	EP 0 548 559 A (AMERICAN CYANAMID CO) 30 June 1993 see pages 3-4, diagrams I, III ----	14-18, 20-32
A	R.P. THUMMEL ET AL.: JOURNAL OF ORGANIC CHEMISTRY, vol. 42, no. 16, 1977, pages 2742-2447, XP002094916 EASTON US see page 2742 ----- -/--	14-18, 20-32

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 February 1999

Date of mailing of the international search report

22/03/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/19788

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 24, no. 1, 1976, pages 85-91, XP002094917 TOKYO JP see chart 1, XI to XII ---	14-18, 20-32
A	DE 36 34 259 A (BASF A.G.) 21 April 1988 see claims ---	14-18, 20-32
A	EP 0 075 727 A (LONZA AG) 6 April 1983 see claims ---	14-18, 20-32
X	CHEMICAL ABSTRACTS, vol. 74, no. 17, 26 April 1971 Columbus, Ohio, US; abstract no. 87279, XP002094918 see Registry Numbers 30989-82-3, 30989-83-4, 30989-84-5, 30989-85-6, 30989-86-7 & E. BREITMAIER ET AL.: CHEM. BER., vol. 104, no. 2, 1971, pages 665-667, ---	33
X	CHEMICAL ABSTRACTS, vol. 91, no. 23, 3 December 1979 Columbus, Ohio, US; abstract no. 193249, XP002094919 see Registry Numbers 71637-33-7, 71637-36-0 & S. GRONOWITZ ET AL.: CHEM. SCR., vol. 13, no. 1, 1979, pages 39-45, ---	33
X	CHEMICAL ABSTRACTS, vol. 92, no. 21, 26 May 1980 Columbus, Ohio, US; abstract no. 180583, XP002094920 see Registry Numbers 73405-93-3, 73405-94-4 & C. SKOETSCH ET AL.: CHEM. BER., vol. 113, no. 2, 1980, pages 795-799, ---	33
X	CHEMICAL ABSTRACTS, vol. 96, no. 25, 21 June 1982 Columbus, Ohio, US; abstract no. 217817, XP002094921 see Registry Number 81927-49-3 & R. HANKE ET AL.: CHEM. BER., vol. 115, no. 4, 1982, pages 1657-1661, ---	33

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/19788

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 99, no. 19, 7 November 1983 Columbus, Ohio, US; abstract no. 158202, XP002094922 see Registry Number 87386-45-6 & L.F. TIETZE ET AL.: TETRAHEDRON LETTERS, vol. 24, no. 34, 1983, pages 3579-3582, OXFORD GB	33
X	--- CHEMICAL ABSTRACTS, vol. 101, no. 13, 24 September 1984 Columbus, Ohio, US; abstract no. 109885, XP002094923 see Registry Numbers 91752-75-9, 91752-76-0 & L. KANIA ET AL.: J. MOL. STRUCT., vol. 117, no. 1-2, 1984, pages 19-31,	33
X	--- CHEMICAL ABSTRACTS, vol. 103, no. 13, 30 September 1985 Columbus, Ohio, US; abstract no. 104142, XP002094924 see Registry Number 97988-75-5 & W. FABIAN: THEOCHEM, vol. 22, 1985, pages 287-297,	33
X	--- CHEMICAL ABSTRACTS, vol. 110, no. 9, 27 February 1989 Columbus, Ohio, US; abstract no. 74786, XP002094925 see Registry Number 116952-07-9 & L.F. TIETZE ET AL.: CHEM. BER., vol. 122, no. 1, 1989, pages 83-94,	33
X	--- CHEMICAL ABSTRACTS, vol. 115, no. 7, 19 August 1991 Columbus, Ohio, US; abstract no. 70882, XP002094926 see Registry Number 135304-92-6 & Z. ARNOLD ET AL.: COLLECT. CZECH. CHEM. COMMUN., vol. 56, no. 5, 1991, pages 1019-1031, --- -/--	33

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 98/19788

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 123, no. 17, 23 October 1995 Columbus, Ohio, US; abstract no. 228067, XP002094927 see Registry Numbers 168145-51-5, 168145-52-6 & S. LINSTROEM: ACTA CHEM. SCAND., vol. 49, no. 5, 1995, pages 361-363, ---	33
X	CHEMICAL ABSTRACTS, vol. 115, no. 13, 30 September 1991 Columbus, Ohio, US; abstract no. 135894, XP002094928 see Registry Number 135987-12-1 & S. KAJIGAESHI ET AL.: CHEM. EXPRESS , vol. 6, no. 7, 1991, pages 527-530, ---	33
X	US 3 277 103 A (S. TROFIMENKO) 4 October 1966 see examples 1,2 ---	33
X	EP 0 017 438 A (AMERICAN CYANAMID CO) 15 October 1980 see page 6, compounds (XV) -----	33

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 98/19788

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9624584 A	15-08-1996	US 5686470 A AU 4859396 A EP 0808304 A	11-11-1997 27-08-1996 26-11-1997
EP 0548559 A	30-06-1993	BR 9205166 A CA 2086281 A JP 5255259 A	29-06-1993 28-06-1993 05-10-1993
DE 3634259 A	21-04-1988	DE 3752066 D EP 0263464 A JP 1090171 A JP 7107050 B US 5079367 A	19-06-1997 13-04-1988 06-04-1989 15-11-1995 07-01-1992
EP 0075727 A	06-04-1983	CH 660733 A AT 24717 T DK 425282 A,B, IE 54022 B US 4421921 A	15-06-1987 15-01-1987 30-03-1983 24-05-1989 20-12-1983
US 3277103 A	04-10-1966	NONE	
EP 0017438 A	15-10-1980	US 4242515 A AT 2674 T AU 533295 B AU 5519680 A CA 1146554 A DD 150466 A JP 55130977 A	30-12-1980 15-03-1986 17-11-1983 02-10-1980 17-05-1983 02-09-1981 11-10-1980